

Simulations of the Discovery of Centaurs and Kuiper Belt Objects

Luke Dones (SJSU Foundation and NASA Ames)

We have performed Monte Carlo simulations of the discovery of Centaurs and Kuiper Belt objects (KBO's) in ecliptic surveys. Our goal is to constrain the populations of these objects and their distributions of size and distance from the Sun. In particular, we apply our model to the new Mauna Kea survey by Jewitt *et al.* (*Astron J.*, in press, Sep. 1996), which discovered 12 KBO's and 2 Centaurs (1994 TA and 1995 DW₂). Because Centaurs typically survive only a few Myr before being ejected from the solar system by one of the giant planets, their numbers should be small compared with the number of KBO's; thus discovery of even one Centaur in a pencil-beam survey is surprising if Centaurs originate from the Kuiper Belt (cf. Irwin *et al.*, *Astron J.* **110**, 3082, 1995). We assume the following: (1) The Kuiper Belt has two components: non-resonant objects with semi-major axes $a > 42$ AU, and objects in the 2:3 resonance at $a = 39$ AU and (possibly) in the 1:2 resonance at $a = 48$ AU (Malhotra, *Astron J.* **110**, 420, 1995). (2) The number of Centaurs increases with distance from the Sun, r , for $10 \text{ AU} \leq r \leq 30 \text{ AU}$, based upon orbital integrations by Levison and Duncan (preprint, 1996). (3) The distribution of diameters is a power law, with differential index q , and an upper cutoff. If we assume $q \sim 3$, the ratio of the observed number of Centaurs to Kuiper Belt objects exceeds the model's predictions by a factor of 10–100. (The large range primarily reflects the small number statistics of the Centaur discoveries.) If we use the steeper size distribution assumed by Weissman and Levison (in *Pluto-Charon*, U. Arizona Press, 1996), which has $q = 4.5$ at large sizes, our model is marginally consistent with the number of objects discovered by Jewitt *et al.* However, if the size distribution were really this steep, the discovered objects would typically be smaller and closer than 1994 TA and 1995 DW₂. On this basis, all models with $q \geq 4$ can be rejected at the 95% confidence level, even allowing for generous uncertainties in the limiting magnitude of the survey and the spatial distribution of the Centaurs. We will present the implications of these results for the origin of the Centaurs.

Abstract submitted for 1996 DPS meeting

Date submitted: LPI electronic form version 5/96

Division for Planetary Sciences Abstract Form

DPS Category 23

Running #7442

Session 0.00

Invited ☐ Poster presentation ☒ Title only ☐

Have you received your Ph.D. since the last DPS meeting?

Yes ☐ No ☐

Is your abstract newsworthy, and if so, would you be willing to prepare a news release and be available for interviews with reporters?

Yes ☐ No ☐ Maybe ☒

Paper presented by Luke Dones

NASA Ames 245-3

Moffett Field CA 94035-1000 USA

Phone: 415-604-0125

Fax: 415-604-6779

Email: luke@boombox.arc.nasa.gov

Special instructions: Tue Aug 27 16:18:12 CDT 1996

Membership Status (First Author):

DPS-AAS Member ☒ Non-Member ☐

Student Member ☐ Student Non-Member ☐

Is this your first DPS presentation? Yes ☐ No ☐

Sponsor: